

www.elsevier.com/locate/jbi




Fig. 1




Fig. 2a

Fig. 2b




Fig. 3a

Fig. 3b

Fig. 3c




Fig. 4




Fig. 5a




Fig. 5b

S# 550-554 RT: 17.50 -17.58AV : 5 NL: 2.02 E8

T: + cFull ms [450.00 - 2000.00]




Fig. 6a

S# 577-596 RT: 18.14-18.59AV:20 NL: 5.72 E6
T: + cFull ms [450.00 - 2000.00]




Fig. 6b

S# 577-596 RT: 18.14-18.59AV:20 NL: 6.04 E7

T: += cFull ms [450.00 - 2000.00]

44600.0




Fig. 6c




Fig. 7a




Fig. 7b




Fig. 7c




Fig. 8a





Fig. 8b


12/14

ScMHC-BirA-Biotin/209



10 $\mu\text{g}/\text{ml}$

Fig. 8c



1 $\mu\text{g}/\text{ml}$


Fig. 8d



0.05 $\mu\text{g}/\text{ml}$

Fig. 8e

CD8-FITC



CD8-FITC

Fig. 9a
Fig. 9b

© 2002 Blackwell Science Ltd, *Journal of Clinical Pharmacy and Therapeutics*, 27, 333–340




Fig. 10a

Fig. 10b